



## Iterative methods for solving split feasibility problems and fixed point problems in Banach spaces

Min Liu<sup>a</sup>, Shi-sen Chang<sup>b</sup>, Ping Zuo<sup>a</sup>, Xiaorong Li<sup>a</sup>

<sup>a</sup>Department of Mathematics, Yibin university, Yibin, Sichuan 644000, China

<sup>b</sup>Center for General Education, China Medical University, Taichung, 40402, Taiwan

### 1 Introduction

Many applications of the split feasibility problem (SFP), which was first introduced by Censor and Elfving [1], have appeared in various fields of science and technology, such as in signal processing, medical image reconstruction and intensity-modulated radiation therapy (for more information, see [2,3] and the references therein). In fact, Censor and Elfving [1] studied SFP in a finite-dimensional space, by considering the problem of finding a point

$$x^* \in C \text{ such that } Ax^* \in Q \quad (1.1)$$

where  $C$  and  $Q$  are nonempty closed convex subsets of  $\mathbb{R}^n$ , and  $A$  is an  $n \times n$  matrix. They introduced an iterative method for solving (SFP) (1.1).

On the other hand, variational inclusion problems are being used as mathematical programming models to study a large number of optimization problems arising in finance, economics, network, transportation and engineering science. The formal form of a variational inclusion problem is to find  $x^* \in H$  such that

$$0 \in Bx^* \quad (1.2)$$

where  $B : H \rightarrow 2^H$  is a set-valued operator. If  $B$  is a maximal monotone operator, the elements in the solution set of the problem (1.2) are called the zero of maximal monotone operator. This problem was introduced by Martinet [4], and later it has been studied by many authors.

It is well known that the popular iteration method that was used for solving the problem (1.2) is the following proximal point algorithm: for a given  $x \in H$ ,

$$x_{n+1} = J_{\lambda_n}^B x_n, \forall n \in \mathbb{N},$$

---

2010 *Mathematics Subject Classification.* 25A18, 47H04, 47H05, 47H10.

*Keywords.* Split feasibility problem; Maximal monotone operators; Shrinking projective method; Modified proximal point algorithm; Fixed point problems; Strong convergence theorem.

Received: 14 August 2019; Accepted: 21 September 2019

Communicated by Dragan S. Djordjević

The corresponding author: Shih-sen Chang

*Email addresses:* liuminybsc@163.com (Min Liu), changss2013@163.com (Shi-sen Chang), zuo1968ping@163.com (Ping Zuo), lixr123456@163.com (Xiaorong Li)

where  $\{\lambda_n\} \subset (0, \infty)$  and  $J_{\lambda_n}^B = (I + \lambda_n B)^{-1}$  is the resolvent of the considered maximal monotone operator  $B$  corresponding to  $\lambda_n$  (see, also [5 – 9]) for more details).

A related topic to the above variational inclusion problem is fixed point theory which has been a very powerful and important tool in the study of mathematical models. Of course, many authors were interested in and studied the approximating of a fixed point of nonlinear mappings by using iterative methods, and applied the obtained results to many important problems, such as the null point problem, variational inequality problem, optimization problems.

For solving the (SFP) and the fixed point problem, Takahashi et al. [10] considered the problem of finding a point  $x^* \in H$  such that

$$0 \in Bx^*, \text{ and } Lx^* \in \text{Fix}(T) \tag{1.3}$$

where  $B : H_1 \rightarrow 2^{H_1}$  is a maximal monotone operator,  $L : H_1 \rightarrow H_2$  is a bounded linear operator and  $T : H_2 \rightarrow H_2$  is a nonexpansive mapping. They considered the following iterative algorithm: for any  $x_1 \in H_1$ ,

$$x_{n+1} = J_{\lambda_n}^B (I - \gamma_n L^* (I - T) L x_n), n \geq 1, \tag{1.4}$$

where  $\{\lambda_n\}$  and  $\{\gamma_n\}$  satisfy some suitable conditions, and  $J_{\lambda_n}^B$  is the resolvent of a maximal monotone operator  $B$  associated to  $\{\lambda_n\}$ . They proved that the sequence  $\{x_n\}$  generated by (1.4) converges weakly to a point  $x^* \in B^{-1}(0) \cap L^{-1}\text{Fix}(T)$ .

Motivated and inspired by [10], the purpose of this paper is to continue to study the problem (1.3) in Banach space. By using shrinking projective method and the modified proximal point algorithm, we propose an iterative algorithm. Under suitable conditions, some strong convergence theorems are proved. Our results extend the corresponding results in Takahashi et al. [10] from Hilbert spaces to Banach spaces. Moreover, the method of proof adopted in this paper is different from that one in [10].

## 2 Preliminaries

Throughout this article, we assume that the Banach space are real. We denote by  $\mathbb{N}$  and  $\mathbb{R}$  the sets of positive integers and real numbers, respectively. Let  $E$  be a Banach space and  $E^*$  be the topological dual of  $E$ . For all  $x \in E$  and  $x^* \in E^*$ , we denote by  $\langle x, x^* \rangle$  the value of  $x^*$  at  $x$ . The mapping  $J : E \rightarrow 2^{E^*}$  defined by

$$J(x) = \{x^* \in E^* : \langle x, x^* \rangle = \|x\|^2 = \|x^*\|^2\}, x \in E \tag{2.1}$$

is called the normalized duality mapping.

A Banach space  $E$  is said to be strictly convex if  $\frac{\|x+y\|}{2} \leq 1$  for all  $x, y \in S(E) = \{z \in E : \|z\| = 1\}$  with  $x \neq y$ . The modulus of convexity of  $E$  is defined by

$$\delta_E(\epsilon) = \inf\{1 - \|\frac{1}{2}(x + y)\| : \|x\| \leq 1, \|y\| \leq 1, \|x - y\| \geq \epsilon\}, \tag{2.2}$$

for all  $\epsilon \in [0, 2]$ .  $E$  is said to be uniformly convex if  $\delta_E(0) = 0$ , and  $\delta_E(\epsilon) > 0$  for all  $0 < \epsilon \leq 2$ . A Hilbert space is 2-uniformly convex.

The modulus of smoothness of  $E$ :  $\rho_E : [0, \infty) \rightarrow [0, \infty)$  is defined by

$$\rho_E(t) = \sup\{\frac{1}{2}(\|x + y\| + \|x - y\|) - 1 : x \in U, \|y\| \leq t\}. \tag{2.3}$$

A Banach space  $E$  is said to be uniformly smooth if  $\frac{\rho_E(t)}{t} \rightarrow 0$  as  $t \rightarrow 0$ . A typical example of uniformly smooth Banach space is  $L^p$ , where  $p > 1$ . More precisely,  $L^p$  is  $\min\{p, 2\}$ -uniformly smooth for every  $p > 1$ . Let  $q$  be a fixed real number with  $q > 1$ , then a Banach space  $E$  is said to be  $q$ -uniformly smooth if there exists a constant  $c > 0$  such that  $\rho_E(t) \leq ct^q$  for all  $t > 0$ . It is well known that every  $q$ -uniformly smooth Banach space is uniformly smooth.

It is also well known that if  $E$  is a uniformly smooth Banach space, then  $J$  is uniformly continuous from norm to norm on each bounded subset of  $E$ .

Let  $C$  be a nonempty, closed, and convex subset of a strictly convex and reflexive Banach  $E$ , then the metric projection

$$P_C x = \arg \min_{y \in C} \|x - y\|, x \in E, \tag{2.4}$$

is the unique minimizer of the norm distance.

Let  $E$  be a smooth, reflexive, and strictly convex Banach space. Consider the functional defined by [11, 12]

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \forall x, y \in E, \tag{2.5}$$

where  $J$  is the normalized duality mapping. It is clear that in a Hilbert space  $H$ , (2.5) reduces to  $\phi(x, y) = \|x - y\|^2, \forall x, y \in H$ .

It is obvious from the definition of  $\phi$  that

$$(\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|x\| + \|y\|)^2, \forall x, y \in E, \tag{2.6}$$

and

$$\phi(x, J^{-1}(\lambda Jy + (1 - \lambda)Jz)) \leq \lambda\phi(x, y) + (1 - \lambda)\phi(x, z), \forall x, y \in E. \tag{2.7}$$

Following Alber [11], the generalized projection  $\Pi_C : E \rightarrow C$  is defined by

$$\Pi_C x = \arg \min_{y \in C} \phi(y, x), \forall x \in E, \tag{2.8}$$

That is,  $\Pi_C(x) = \bar{x}$ , where  $\bar{x}$  is the unique solution to the minimization problem  $\phi(\bar{x}, x) = \inf_{y \in C} \phi(y, x)$ . The existence and uniqueness of the operator  $\Pi_C$  follows from the properties of the functional  $\phi(x, y)$  and strict monotonicity of the mapping  $J$ (see, e.g., [11 – 14]). In Hilbert space  $H, \Pi_C = P_C$ .

Let  $T : E \rightarrow E$  be a mapping. we say that  $T$  is nonexpansive, if

$$\|Tx - Ty\| \leq \|x - y\|, \forall x, y \in E. \tag{2.9}$$

A set-valued mapping  $A : E \rightarrow 2^E$  is said to be monotone if for any  $x, y \in E$  and any  $x^* \in Ax, y^* \in Ay$ , we have

$$\langle x^* - y^*, x - y \rangle \geq 0. \tag{2.10}$$

A monotone operator  $A$  is said to be maximal if its  $Gr(A) = \{(x, x^*) : x^* \in Ax\}$  is not properly contained in graph of any other monotone operator.

If  $E$  is a strictly convex, reflexive, and smooth Banach space, and  $B : E \rightarrow 2^E$  is a maximal monotone operator then, for any positive real number  $\lambda$ , we can define a single-valued mapping  $J_\lambda^B : E \rightarrow E$  by

$$J_\lambda^B(x) = (J + \lambda B)^{-1}J(x), x \in E. \tag{2.11}$$

This mapping is called the resolvent of  $B$  for  $\lambda > 0$ .

It is know that  $B^{-1}(0) = F(J_\lambda^B)$  for every  $\lambda > 0$  and  $B^{-1}(0)$  is a closed and convex subset of  $E$ . For more details, see [15-17].

**Lemma 2.1** [18] Let  $E$  be a strictly convex and reflexive smooth Banach space. Let  $B : E \rightarrow 2^E$  be a maximal monotone operator and  $J_\lambda^B$  be the resolvent of  $B$  for  $\lambda > 0$ , then

$$\phi(u, J_\lambda^B x) \leq \phi(u, x)$$

for any  $u \in B^{-1}(0)$  and  $x \in E$ .

**Lemma 2.2**[14] Let  $E$  be a uniformly convex and smooth Banach space and let  $\{x_n\}$  and  $\{y_n\}$  be two sequences of  $E$ . If  $\phi(x_n, y_n) \rightarrow 0$  and either  $\{x_n\}$  or  $\{y_n\}$  is bounded, then  $\|x_n - y_n\| \rightarrow 0$ .

**Lemma 2.3**<sup>[12]</sup> Let  $E$  be a smooth, strictly convex, and reflective Banach space and  $C$  be nonempty closed convex subset of  $E$ . Then, the following conclusions hold:

- (1)  $\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y), \forall x \in C, y \in E$ .
- (2) If  $x \in E$  and  $z \in C$ , then  $z = \Pi_C x$  iff  $\langle z - y, Jx - Jz \rangle \geq 0, \forall y \in C$ .
- (3) For  $x, y \in E, \phi(x, y) = 0$  if and only if  $x = y$ .

**Lemma 2.4**<sup>[17]</sup> Let  $E$  be a smooth, strictly convex, and reflective Banach space. Let  $C$  be a nonempty, closed, and convex subset of  $E$  and let  $x_1 \in E$  and  $z \in C$ . Then, the following conclusions hold:

- (1)  $z = P_C x_1$ .
- (2)  $\langle z - y, J(x_1 - z) \rangle \geq 0, \forall y \in C$ .

**Lemma 2.5**<sup>[19]</sup> For a given number  $r > 0$ , a real Banach space  $E$  is uniformly convex if and only if there exists a continuous strictly increasing function  $g : [0, \infty) \rightarrow [0, \infty)$  with  $g(0) = 0$  such that

$$\|tx + (1 - t)y\|^2 \leq t\|x\|^2 + (1 - t)\|y\|^2 - t(1 - t)g\|x - y\|,$$

for all  $x, y \in E$  with  $\|x\| \leq r$  and  $\|y\| \leq r$ , and  $t \in [0, 1]$

**Lemma 2.6**<sup>[19]</sup> Let  $E$  be a 2-uniformly smooth Banach space with the best smoothness constants  $\kappa > 0$ . Then, the following inequality holds:

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, Jx \rangle + 2\|\kappa y\|^2, \forall x, y \in E.$$

### 3 The Main Results

**Theorem 3.1** Let  $E_1$  be a real 2-uniformly convex and 2-uniformly smooth Banach space with the best smoothness constant  $\kappa > 0$ . Let  $E_2$  be a smooth strictly convex and reflective Banach space. Let  $B : E_1 \rightarrow 2^{E_1}$  be a maximal monotone operator,  $L : E_1 \rightarrow E_2$  be a bounded linear operator with adjoint  $L^*$ , and  $T : E_2 \rightarrow E_2$  be a nonexpansive mappings. Let  $x_1 \in E_1, C_1 = E_1$  and  $\{x_n\}$  be a sequence generated by

$$\begin{cases} z_n = J_1^{-1}(J_1 x_n + \gamma L^* J_2 (P_{F(T)} - I) L x_n), n \geq 1, \\ y_n = J_1^{-1}[(1 - \alpha_n) J_1 z_n + \alpha_n J_1 J_\lambda^B z_n], \\ C_{n+1} = \{v \in C_n : \phi(v, y_n) \leq \phi(v, x_n); \phi(v, z_n) \leq \phi(v, x_n)\}, \\ x_{n+1} = \Pi_{C_{n+1}} x_1, \end{cases} \quad (3.1)$$

where  $J_\lambda^B = (J_1 + \lambda B)^{-1} J_1$ . If  $Q = \{p \in B^{-1}(0) : Lp \in F(T)\} \neq \emptyset$  and the following conditions are satisfied:

- (1)  $\alpha_n \in [\delta, 1), \delta > 0$ ,
- (2)  $0 < \gamma < \frac{1}{\kappa^2 \|L\|^2}$ ,

then  $\lim_{n \rightarrow \infty} x_n = x^* = \Pi_Q x_1$  which is a solution of problem (1.3).

**Proof** We shall divide the proof into five steps.

**Step 1** First we show that  $C_n$  is a closed and convex subset for each  $n \geq 1$ .

Since  $C_1 = E_1, C_1$  is closed and convex. Assume that  $C_n$  is closed and convex for some  $n \geq 2$ . Next, by induction, we prove that  $C_{n+1}$  is also closed and convex. In fact, it follows from (3.1) that for any  $v \in C_n$ , we have

$$\phi(v, y_n) \leq \phi(v, x_n) \iff 2\langle v, Jx_n - Jy_n \rangle \leq \|x_n\|^2 - \|y_n\|^2, \quad (3.2)$$

$$\phi(v, z_n) \leq \phi(v, x_n) \iff 2\langle v, Jx_n - Jz_n \rangle \leq \|x_n\|^2 - \|z_n\|^2. \quad (3.3)$$

These imply that  $C_{n+1}$  is a closed and convex subset of  $E_1$ .

**Step 2** Now we prove that  $Q \subseteq C_n$ , for all  $n \geq 1$ .

Let  $p \in Q$ . By the assumption that  $E_1$  is a 2-uniformly convex and 2-uniformly smooth real Banach space, so,  $E_1^*$  is 2-uniformly smooth real Banach space and  $J_1 = (J_1^*)^{-1}$ . It follows from (3.1) and Lemma 2.6 that

$$\begin{aligned}
 \phi(p, z_n) &= \phi(p, J_1^{-1}(J_1x_n + \gamma L^* J_2(P_{F(T)} - I)Lx_n)) \\
 &= \|p\|^2 - 2\langle p, J_1x_n + \gamma L^* J_2(P_{F(T)} - I)Lx_n \rangle \\
 &\quad + \|J_1x_n + \gamma L^* J_2(P_{F(T)} - I)Lx_n\|^2 \\
 &\leq \|p\|^2 - 2\langle p, J_1x_n \rangle - 2\langle p, \gamma L^* J_2(P_{F(T)} - I)Lx_n \rangle + \|x_n\|^2 \\
 &\quad + 2\gamma \langle Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle + 2k^2 \|L\|^2 \gamma^2 \|(P_{F(T)} - I)Lx_n\|^2 \\
 &\leq \phi(p, x_n) - 2\gamma \langle Lp - Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle \\
 &\quad + 2k^2 \|L\|^2 \gamma^2 \|(P_{F(T)} - I)Lx_n\|^2.
 \end{aligned} \tag{3.4}$$

Furthermore from Lemma 2.4 we have that

$$\begin{aligned}
 \langle Lp - Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle &= \langle Lp - P_{F(T)}Lx_n + P_{F(T)}Lx_n - Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle \\
 &= \langle Lp - P_{F(T)}Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle \\
 &\quad + \langle (P_{F(T)} - I)Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle \\
 &= \langle Lp - P_{F(T)}Lx_n, J_2(P_{F(T)} - I)Lx_n \rangle + \|(P_{F(T)} - I)Lx_n\|^2 \\
 &\geq \|(P_{F(T)} - I)Lx_n\|^2.
 \end{aligned} \tag{3.5}$$

Substituting (3.5) into (3.4), and by using the condition (2) to simplify, we have

$$\phi(p, z_n) \leq \phi(p, x_n) - 2\gamma(1 - k^2 \|L\|^2 \gamma) \|(P_{F(T)} - I)Lx_n\|^2 \leq \phi(p, x_n). \tag{3.6}$$

Furthermore, from Lemma 2.1, Lemma 2.5, (3.1) and (3.6), we obtain

$$\begin{aligned}
 \phi(p, y_n) &= \phi(p, J_1^{-1}[(1 - \alpha_n)J_1z_n + \alpha_n J_1 J_\lambda^B z_n]) \\
 &= \|p\|^2 - 2\langle p, (1 - \alpha_n)J_1z_n + \alpha_n J_1 J_\lambda^B z_n \rangle \\
 &\quad + \|(1 - \alpha_n)J_1z_n + \alpha_n J_1 J_\lambda^B z_n\|^2 \\
 &\leq \|p\|^2 - 2(1 - \alpha_n)\langle p, J_1z_n \rangle - 2\alpha_n \langle p, J_1 J_\lambda^B z_n \rangle \\
 &\quad + (1 - \alpha_n)\|z_n\|^2 + \alpha_n \|J_\lambda^B z_n\|^2 - \alpha_n(1 - \alpha_n)g(\|J_1z_n - J_1 J_\lambda^B z_n\|) \\
 &= (1 - \alpha_n)\phi(p, z_n) + \alpha_n \phi(p, J_\lambda^B z_n) \\
 &\quad - \alpha_n(1 - \alpha_n)g(\|J_1z_n - J_1 J_\lambda^B z_n\|) \\
 &\leq (1 - \alpha_n)\phi(p, z_n) + \alpha_n \phi(p, z_n) - \alpha_n(1 - \alpha_n)g(\|J_1z_n - J_1 J_\lambda^B z_n\|) \\
 &\leq (1 - \alpha_n)\phi(p, x_n) + \alpha_n \phi(p, x_n) - \alpha_n(1 - \alpha_n)g(\|J_1z_n - J_1 J_\lambda^B z_n\|) \\
 &\leq \phi(p, x_n).
 \end{aligned} \tag{3.7}$$

It follows from (3.6) and (3.7) that  $p \in C_{n+1}$ . This implies that  $Q \subseteq C_n$  for all  $n \geq 1$ .

Therefore,  $\Pi_{C_{n+1}}x_1$  is well defined.

**Step 3** Now we prove that  $\{x_n\}$  is a Cauchy sequence.

Let  $p \in Q$ , by the definition of  $C_n$ , We have  $x_n = \Pi_{C_n}x_1$  for all  $n \geq 1$ . Hence It follows from Lemma 2.3 that

$$\phi(x_n, x_1) = \phi(\Pi_{C_n}x_1, x_1) \leq \phi(p, x_1) - \phi(p, \Pi_{C_n}x_1) \leq \phi(p, x_1), \forall n \geq 1. \tag{3.8}$$

This implies that  $\{\phi(x_n, x_1)\}$  is bounded. In addition, since  $x_n = \Pi_{C_n}x_1$  and

$$x_{n+1} = \Pi_{C_{n+1}}x_1 \in C_{n+1} \subseteq C_n,$$

we have

$$\phi(x_n, x_1) \leq \phi(x_{n+1}, x_1), \forall n \geq 1. \tag{3.9}$$

Therefore,  $\{\phi(x_n, x_1)\}$  is nondecreasing and bounded. So the limit  $\lim_{n \rightarrow \infty} \phi(x_n, x_1)$  exists. Hence from Lemma 2.3, we have

$$\phi(x_{n+1}, x_n) = \phi(x_{n+1}, \Pi_{C_n}x_1) \leq \phi(x_{n+1}, x_1) - \phi(\Pi_{C_n}x_1, x_1) = \phi(x_{n+1}, x_1) - \phi(x_n, x_1), \tag{3.10}$$

which implies that

$$\lim_{n \rightarrow \infty} \phi(x_{n+1}, x_n) = 0. \tag{3.11}$$

This together with Lemma 2.2 shows that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \tag{3.12}$$

For some positive integers  $m, n$  with  $m \leq n$ , it follows from  $x_n = \prod_{C_n} x_1 \subseteq C_m$  and Lemma 2.2 that

$$\begin{aligned} \phi(x_m, x_n) &= \phi(x_m, \prod_{C_n} x_1) \\ &\leq \phi(x_m, x_1) - \phi(\prod_{C_n} x_1, x_1) \\ &= \phi(x_m, x_1) - \phi(x_n, x_1). \end{aligned} \tag{3.13}$$

Since  $\lim_{n \rightarrow \infty} \phi(x_n, x_1)$  exists, it follows from (3.13) and Lemma 2.2 that  $\lim_{n \rightarrow \infty} \|x_n - x_m\| = 0$ . Therefore,  $\{x_n\}$  is a cauchy sequence.

**Step 4** Now we prove that  $\lim_{n \rightarrow \infty} \|z_n - J_\lambda^B z_n\| = 0$  and  $\lim_{n \rightarrow \infty} \|(P_{F(T)} - I)Lx_n\| = 0$ .  
 Since  $x_{n+1} = \prod_{C_{n+1}} x_1 \in C_{n+1} \subseteq C_n$ , by the definition of  $C_{n+1}$  and (3.11), we have

$$\begin{aligned} \phi(x_{n+1}, z_n) &\leq \phi(x_{n+1}, x_n) \rightarrow 0 \text{ (as } n \rightarrow \infty), \\ \phi(x_{n+1}, y_n) &\leq \phi(x_{n+1}, x_n) \rightarrow 0, \text{ (as } n \rightarrow \infty). \end{aligned} \tag{3.14}$$

By Lemma 2.2, we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - z_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|x_{n+1} - y_n\| = 0, \tag{3.15}$$

and so

$$\lim_{n \rightarrow \infty} \|y_n - z_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|x_n - z_n\| = 0. \tag{3.16}$$

Furthermore, it follows from (3.6) that

$$\begin{aligned} 2\gamma(1 - k^2\|L\|^2\gamma)\|(P_{F(T)} - I)Lx_n\|^2 &\leq \phi(p, x_n) - \phi(p, z_n) \\ &= \|p\|^2 - 2\langle p, J_1 x_n \rangle + \|x_n\|^2 - \|p\|^2 + 2\langle p, J_1 z_n \rangle - \|z_n\|^2 \\ &= 2\langle p, J_1 z_n - J_1 x_n \rangle + \|x_n\|^2 - \|z_n\|^2 \\ &\leq 2\|p\| \cdot \|J_1 z_n - J_1 x_n\| + \|x_n - z_n\| \cdot (\|x_n\| + \|z_n\|). \end{aligned} \tag{3.17}$$

Since  $E_1$  is a 2-uniformly convex and 2-uniformly smooth real Banach space,  $J_1$  is uniformly continuous from norm-to-norm. By (3.16), we have  $\|J_1 z_n - J_1 x_n\| \rightarrow 0$ . Also, from (3.17) and the condition (2), we have

$$\lim_{n \rightarrow \infty} \|(P_{F(T)} - I)L(x_n)\| = 0. \tag{3.18}$$

Again from (3.1) we have

$$\alpha_n \|J_1 J_\lambda^B z_n - J_1 z_n\| = \|J_1 y_n - J_1 z_n\|. \tag{3.19}$$

Hence from (3.16) and condition (1) we have

$$\lim_{n \rightarrow \infty} \|J_1 J_\lambda^B z_n - J_1 z_n\| = 0, \tag{3.20}$$

which implies that

$$\lim_{n \rightarrow \infty} \|J_\lambda^B z_n - z_n\| = 0. \tag{3.21}$$

**Step 5** Now we prove that  $\lim_{n \rightarrow \infty} x_n = x^* = \Pi_Q x_1$ .

In fact, since  $\{x_n\}$  is a cauchy sequence, there exists  $x^* \in E_1$  such that  $\lim_{n \rightarrow \infty} x_n = x^*$ . It follows from (3.16) and (3.21) that

$$\lim_{n \rightarrow \infty} z_n = x^*, \lim_{n \rightarrow \infty} J_\lambda^B z_n = x^*. \tag{3.22}$$

Since  $J_\lambda^B = (J_1 + \lambda B)^{-1}J_1$ , we have that  $\frac{J_1 z_n - J_1 J_\lambda^B z_n}{\lambda} \in B J_\lambda^B z_n$ , for all  $n \in \mathbb{N}$ . From the monotonicity of  $B$ , we have that

$$0 \leq \langle u - J_\lambda z_n, v^* - \frac{J_1 z_n - J_1 J_\lambda z_n}{\lambda} \rangle,$$

for all  $(u, v^*) \in Gr(B)$ . Taking  $n \rightarrow \infty$ , we have from (3.21) and (3.22) that  $0 \leq \langle u - x^*, v^* - 0 \rangle$  for all  $(u, v^*) \in Gr(B)$ . Since  $B$  is maximal monotone, we have  $x^* \in B^{-1}(0)$ . In addition, from Lemma 2.4, we obtain

$$\begin{aligned} \|(I - P_{F(T)})Lx^*\|^2 &= \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx^* - P_{F(T)}(Lx^*) \rangle \\ &= \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx^* - Lx_n + Lx_n \\ &\quad - P_{F(T)}(Lx_n) + P_{F(T)}(Lx_n) - P_{F(T)}(Lx^*) \rangle \\ &= \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx^* - Lx_n \rangle \\ &\quad + \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx_n - P_{F(T)}(Lx_n) \rangle \\ &\quad + \langle J_2(Lx^* - P_{F(T)}(Lx^*)), P_{F(T)}(Lx_n) - P_{F(T)}(Lx^*) \rangle \\ &\leq \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx^* - Lx_n \rangle \\ &\quad + \langle J_2(Lx^* - P_{F(T)}(Lx^*)), Lx_n - P_{F(T)}(Lx_n) \rangle. \end{aligned} \tag{3.23}$$

Since  $L$  is a bounded linear operator, we have that  $\lim_{n \rightarrow \infty} \|Lx_n - Lx^*\| = 0$ . Hence by (3.18) we get  $\|(I - P_{F(T)})(Lx^*)\| = 0$ . This implies that  $Lx^* \in F(T)$ . Therefore we have  $x^* \in Q$ .

Let  $z = \Pi_Q x_1, z \in Q$ . From  $x_n = \Pi_{C_n} x_1$  and  $z \in Q \subseteq C_n$ , we have

$$\phi(x_n, x_1) \leq \phi(z, x_1). \tag{3.24}$$

This implies that

$$\phi(x^*, x_1) \leq \lim_{n \rightarrow \infty} \phi(x_n, x_1) \leq \phi(z, x_1). \tag{3.25}$$

By the definition of  $z = \Pi_Q x_1$ , we have  $x^* = z$ . Therefore,  $\lim_{n \rightarrow \infty} x_n = x^* = \Pi_Q x_1$ .

The proof of Theorem 3.1 is completed.

The following result can be obtained from Theorem 3.1 immediately.

**Corollary 3.2** Let  $H_1$  and  $H_2$  be two real Hilbert space. Let  $B : H_1 \rightarrow 2^{H_1}$  be a maximal monotone operator,  $L : H_1 \rightarrow H_2$  be a bounded linear operator with adjoint  $L^*$ ,  $T : H_2 \rightarrow H_2$  be a nonexpansive mapping.

Let  $x_1 \in H$  and  $C_1 = H_1$ , and  $\{x_n\}$  be a sequence generated by

$$\begin{cases} z_n = x_n + \gamma L^*(P_{F(T)} - I)Lx_n, n \geq 1, \\ y_n = (1 - \alpha_n)z_n + \alpha_n J_\lambda^B z_n \\ C_{n+1} = \{v \in C_n : \|y_n - v\| \leq \|x_n - v\|; \|z_n - v\| \leq \|x_n - v\|\}, \\ x_{n+1} = P_{C_{n+1}} x_1, \end{cases} \tag{3.26}$$

where  $P_{F(T)}$  is the metric projection of  $H_2$  onto  $F(T)$  and  $P_{C_{n+1}}$  is the metric projection of  $H_1$  onto  $C_{n+1}$ ,  $J_\lambda^B = (I + \lambda B)^{-1}$ , for  $\lambda > 0$ . If  $Q = \{p \in B^{-1}(0) : Lp \in F(T)\} \neq \emptyset$  and the following conditions are satisfied:

- (1)  $\alpha_n \in [\delta, 1), \delta > 0$ ;
- (2)  $0 < \gamma < \frac{1}{\|L\|^2}$ .

then  $\lim_{n \rightarrow \infty} x_n = x^* = P_Q x_1$ .

**Remark 3.3** Theorem 3.1 generalizes the main results in [10] from Hilbert space to Banach space and the weak convergence of the iterative extends to strong convergence. Moreover, the method of proof is different from that one in [10].

#### 4 Application to split null point problem (SNPP)

Let  $H_1, H_2$  be two real Hilbert spaces. Let  $B_1 : H_1 \rightarrow 2^{H_1}$  and  $B_2 : H_2 \rightarrow 2^{H_2}$  be two set-valued maximal monotone mappings and  $L : H_1 \rightarrow H_2$  be a bounded linear operators. The “so-called” (SNPP) is to find a point  $x^* \in H_1$ , such that

$$0 \in B_1(x^*) \text{ and } 0 \in B_2(Lx^*). \tag{4.1}$$

For solving problem (4.1), Byrne et al. [20] proposed the following iterative algorithm, for  $\lambda > 0$  and an arbitrary  $x_1 \in H_1$ ,

$$x_{n+1} = J_\lambda^{B_1}(x_n - \gamma L^*(I - J_\lambda^{B_2})Lx_n), \forall n \in \mathbb{N}$$

where  $L^*$  is the adjoint of  $L$ ,  $\gamma \in (0, \frac{2}{\|L\|^2})$ ,  $J_\lambda^{B_1}$  and  $J_\lambda^{B_2}$  are the resolvents of maximal monotone operators  $B_1$  and  $B_2$ , respectively. Under suitable control conditions, they proved that  $\{x_n\}$  converges weakly to a point  $x^*$  which is a solution of problem (4.1).

Since  $B_2 : H_2 \rightarrow 2^{H_2}$  is a maximal monotone operator, it is well know that its resolvent operator  $J_\lambda^{B_2} = (I + \lambda B_2)^{-1}$  is nonexpansive and  $B_2^{-1}(0) = F(J_\lambda^{B_2})$ . Then problem (4.1) is equivalent to find a point  $x^* \in H$  such that

$$0 \in B_1(x^*) \text{ and } Lx^* \in B_2^{-1}(0) = F(J_\lambda^{B_2}). \tag{4.2}$$

Then the following result can be obtained from Corollary 3.2 immediately.

**Theorem 4.1** Let  $B_1 : H_1 \rightarrow 2^{H_1}$  and  $B_2 : H_2 \rightarrow 2^{H_2}$  be maximal monotone operators, and  $L : H_1 \rightarrow H_2$  be a bounded linear operator with adjoint  $L^*$ . Let  $x_1 \in H_1$  and  $C_1 = H_1$ , and  $\{x_n\}$  be a sequence generated by

$$\begin{cases} z_n = x_n + \gamma L^*(P_{F(J_\lambda^{B_2})} - I)Lx_n, n \geq 1, \\ y_n = (1 - \alpha_n)z_n + \alpha_n J_\lambda^{B_1} z_n, \\ C_{n+1} = \{v \in C_n : \|y_n - v\| \leq \|x_n - v\|; \|z_n - v\| \leq \|x_n - v\|\}, \\ x_{n+1} = P_{C_{n+1}} x_1, \end{cases} \tag{4.3}$$

where  $P_{F(J_\lambda^{B_2})}$  is the metric projection of  $H_2$  onto  $F(J_\lambda^{B_2})$  and  $P_{C_{n+1}}$  is the metric projection of  $H_1$  onto  $C_{n+1}$ ,  $J_\lambda^{B_1} = (I + \lambda B_1)^{-1}$ ,  $J_\lambda^{B_2} = (I + \lambda B_2)^{-1}$  for  $\lambda > 0$ . If  $Q = \{p \in B_1^{-1}(0) : Lp \in B_2^{-1}(0)\} \neq \emptyset$  and the following conditions are satisfied:

- (1)  $\alpha_n \in [\delta, 1)$ ,  $\delta > 0$ ;
- (2)  $0 < \lambda < \frac{1}{\|L\|^2}$ .

Then the sequence  $\{x_n\}$  converges strongly to a point  $x^* = P_Q x_1$ , which is a solution of problem (4.2).

**Funding** This project is supported by Scientific Research Fund of Science and Technology Department of Sichuan Province (No.2018JY0340). This work was also supported by the Natural Science Foundation of China Medical University, Taiwan.

**Acknowledgements** The authors would to express their thanks to the Editors and the Referees for their helpful com- ments and advices.

**References**

- [1] Censor,Y., Elfving, T.: *A multiprojection algorithm using Bregman projections in product space*, Algorithms 8,221-239 (1994).
- [2] Byrne,C.: *Iterative oblique projection onto convex sets and the split feasibility problem*, Inverse Probl. 18, 441-453 (2002).
- [3] Censor, Y., Bortfeld, T., Martin, B., Trofimov, A.: *A unified approach for inversion problems in intensity-modulated radiation therapy*, Phys. Med. Biol. 51, 2353-2365 (2006).
- [4] Martinet, B.,: *Régularisation d'inéquations variationnelles par approximations successives*, Rev. Fr. Inform. Rech. Opér. 3, 154-158 (1970).
- [5] Bruck,RE, Reich, S.: *Nonexpansive projections and resolvents of accretive operators in Banach space*, Houst. J. Math. 3, 154-158 (1977).
- [6] Eckstein,J., Bertsekas, DP: *On the Douglas Rachford splitting method and proximal point algorithm for maximal monotone operators*, Math. Program. 55, 293-318 (1992).
- [7] Marino,G., Xu,HK: *Convergence of generalized proximal point algorithm*, Commun. Pure Appl. Anal. 3, 791-808 (2004).



- [8] Xu, HK: *Iterative algorithms for nonlinear operators*, J. Lond. Math. Soc. 66, 240-256 (2002).
- [9] Yao, Y., Noor, MA: *On convergence criteria of generalized proximal point algorithms*, J. Comput. Appl. Math. 217, 46-55 (2008).
- [10] Takahashi, W., Xu, HK, Yao, JC: *Iterative methods for generalized split feasibility problems in Hilbert space*, Set-Valued Var. Anal. 23, 205-221 (2015).
- [11] Alber, Ya.I: *Metric and generalized projection operators in Banach space: properties and application*, In; Kartsatos, A.G.(ed.) *Theory and Applications of Nonlinear Operators of Accretive and Monotonic Type*, pp. 15-50. Marcel Dekker, New York (1996).
- [12] Alber, Ya.I., Reich, S.: *An iterative method for solving a class of nonlinear operator equations in Banach space*, Panamer. Math. J. 4(2), 39-54 (1994).
- [13] Cioranescu, I.: *Geometry of Banach space, Duality Mappings and Nonlinear Problems*, Kluwer Academic Publishers, Dordrecht (1990).
- [14] Kamimura, S., Takahashi, W.: *Strong convergence of a proximal-type algorithm in Banach space*, SIAM. J. Optim. 13, 938-945 (2002).
- [15] Barbu, V., Precupanu, T.: *Convexity Optimization in Banach space*, Romanian ed. Math. Appl. (East Eur. Ser.) Vol. 10. d. Reidel publishing Codordrecht (1986).
- [16] Takiahashi, W.: *Convex Analysis and Approximation of Fixed points*, Yokohama Pherubl, Yokohama. 2000 (in Japanese).
- [17] Takiahashi, W. ; *Nonlinear Functional Analysis: Fixed point Theory and Its Applications*, Yokohama publ., Yokohama (2000).
- [18] Matsushita, S., Takahashi, W.: *A strong convergence theorem for relatively nonexpansive mappings in Banach space*, J. Approx. Theory. 134, 257-266 (2005).
- [19] Xu, HK: *Inequalities in Banach spaces with applications*, Nonlinear Anal. Theory Methods Appl. 16, 1127-1138 (1991).
- [20] Byrne, C., Censor, Y., Gibali, A., Reich, S.: *Weak and strong convergence of algorithms for the split common null point problem*, J. Nonlinear Convex Anal. 13, 759-775 (2012).