

## ON THE RATIO OF DIRECTED LENGTHS ON THE PLANE WITH GENERALIZED ABSOLUTE VALUE METRIC AND RELATED PROPERTIES

Gökhan Soydan<sup>1</sup>, Yusuf Dođru<sup>2</sup> and N.Umut Arslandođan<sup>3</sup>

### Abstract

In this paper, we show that a point of division divides a related line segment in the same ratio on the plane with generalized absolute value metric and Euclidean plane. Then the coordinates of the division point can be determined by the same formula as in the Euclidean plane. In the latter parts of the work, we give Ceva's and Menelaus's theorems and the theorem of directed lines on the plane with generalized absolute value metric.

## 1 Introduction

H. Minkowski [22] published a whole family of metrics providing new insight into the study of plane geometry at the beginning of last century. Later, taxicab plane geometry was introduced by Menger, and developed by Krause, using the metric  $d_T(K, L) = |x_1 - x_2| + |y_1 - y_2|$  instead of the well-known Euclidean metric  $d_E(K, L) = ((x_1 - x_2)^2 + (y_1 - y_2)^2)^{1/2}$  for the distance between any two points  $K = (x_1, y_1)$  and  $L = (x_2, y_2)$  in the Cartesian coordinate plane (see [20], [17]). Recently, the plane geometry with the generalized absolute value metric  $d_G$  was introduced by Kaya et al. [13] and Akca et al. [3]. In  $\mathbb{R}^2$ , the  $d_G$ -distances between  $K$  and  $L$  are defined by

$$\begin{aligned} d_G(K, L) &= d_G((x_1, y_1), (x_2, y_2)) \\ &= k_1 \max\{|x_1 - x_2|, |y_1 - y_2|\} + k_2 \min\{|x_1 - x_2|, |y_1 - y_2|\} \end{aligned} \quad (1)$$

for all  $k_1, k_2 \in \mathbb{R}$ ,  $k_1 \geq k_2 \geq 0$ ,  $k_1 \neq 0$ . According to the definition of  $d_G$ -distances, the shortest way between two points  $K$  and  $L$  is the union of a vertical or a horizontal line segment and a line segment with the slope  $\pm 2k_1k_2/(k_1^2 - k_2^2)$ .

---

Corresponding Author<sup>1</sup>

2010 *Mathematics Subject Classifications*. 51K05, 51K99.

*Key words and Phrases*. Ceva theorem, distance, metric, Menelaus theorem, protractor geometry.

Received: November 24, 2009

Communicated by Vladimir Dragovic

The family of distances,  $d_G$ , which includes Chinese checkers, taxicab distance and maximum distance as special cases.

A *metric geometry* consists of a set  $P$  whose elements are called *points*, together with a collection  $L$  of non-empty subsets of  $P$ , called *lines*, and a distance function  $d$ , such that

- 1) Every two distinct points in  $P$  lie on a unique line,
- 2) There exist three points in  $P$ , which do not lie all on one line,
- 3) There exists a bijective function  $f : l \rightarrow \mathbb{R}$  for all lines in  $L$  such that  $|f(P) - f(Q)| = d(P, Q)$  for each pair of points  $P$  and  $Q$  on  $l$ .

A metric geometry defined above is denoted by  $\{P, L, d\}$ . However, if a metric geometry satisfies the plane separation axiom below, and it has an angle measure function  $m$ , then it is called *protractor geometry* and denoted by  $\{P, L, d, m\}$ .

- 4) For every  $l$  in  $L$ , there are two subsets  $H_1$  and  $H_2$  of  $P$  (called *half planes* determined by  $l$ ) such that

- (i)  $H_1 \cup H_2 = P - l$  ( $P$  with  $l$  removed),
- (ii)  $H_1$  and  $H_2$  are disjoint and each is convex,
- (iii) If  $A \in H_1$  and  $B \in H_2$ , then  $[AB] \cap l = \emptyset$ .

If  $L_E$  is the set of all lines in the Cartesian coordinate plane, and  $m_E$  is the standard angle measure function in the Euclidean plane, then  $\{\mathbb{R}^2, L_E, d_C, m_E\}$ , called the plane with the generalized absolute value metric is a model of protractor geometry (This can be shown easily: the proof is similar to that of taxicab plane; refer to [21] or [8] to see that the taxicab plane is a model of protractor geometry). The plane with the generalized absolute value metric is also in the class of non-Euclidean geometries since it fails to satisfy the side-angle-side axiom. However, the plane with the generalized absolute value metric is almost the same as Euclidean plane  $\{\mathbb{R}^2, L_E, d_C, m_E\}$  since the points are the same, the lines are the same and the angles are measured in the same way. Since the plane with the generalized absolute value metric has distance function different from that in the Euclidean plane, it is interesting to study on the plane with the generalized absolute value metric of topics that include the distance concept in the Euclidean plane. ([1], [2], [4], [5], [6], [7], [9], [12], [10], [11], [13], [14], [16], [23], [24], [25], [19], [26], [27], [28], [29]) These topics are division point, directed lengths, ratio of directed lengths, Menelaus' Theorem, Ceva's Theorem and the theorem of directed lines. In this paper, GAM is the abbreviation for the plane geometry with the generalized absolute value metric.

## 2 On The Ratio Of Directed Lengths On The Plane With Generalized Absolute Value Metric

Let  $A$  and  $B$  be any two points on a directed straight line  $l$ . We define *directed* GAM length of the line segment  $AB$  as follows:

$$d_G[AB] = \begin{cases} d_G(A, B), & \text{if } AB \text{ and } l \text{ have the same direction} \\ -d_G(A, B), & \text{if } AB \text{ and } l \text{ have the opposite direction.} \end{cases}$$

Thus,  $d_G[AB] = -d_G[BA]$ . Clearly, *directed length* in the Euclidean plane can be defined in a similar way. That is

$$d_E[AB] = \begin{cases} d_E(A, B), & \text{if } AB \text{ and } l \text{ have the same direction} \\ -d_E(A, B), & \text{if } AB \text{ and } l \text{ have the opposite direction.} \end{cases}$$

If  $P, Q, R$  are points on a same directed line and  $R$  is between points  $P$  and  $Q$ , we denote this by  $PRQ$ . If  $PRQ$ , then  $R$  divides the line segment  $PQ$  *internally* and the ratio of the directed GAM lengths is a positive real number, that is

$d_G[PR]/d_G[RQ] = \delta > 0$ . If  $PQR$  or  $RPQ$  then  $R$  divides the line segments  $PQ$  *externally*,  $d_G[PR]/d_G[RQ] = \delta < 0$ , that is, the line segments  $PR$  and  $RQ$  have opposite directions. In both cases  $R$  called the *division point* which divides the line segment  $PQ$  in ratio  $\delta$ . It's obvious,  $R \neq Q$ .  $R = P \Leftrightarrow \delta = 0$  and ( $R$  is at infinity  $\Leftrightarrow \delta = -1$ ).

Let  $R$  and  $R'$  be two points such that  $R$  divides a given line segment  $PQ$  internally and  $R'$  divides  $PQ$  externally in the same proportion though with opposite signs. Thus, the ratio of the directed lengths,  $d_G[PR]/d_G[RQ] = -d_G[PR']/d_G[R'Q]$  is the same positive number  $\delta$ .

**Lemma 1.** *For any two points  $K$  and  $L$  in the Cartesian plane that do not lie on a vertical line, if  $m$  is the slope of the line through  $K$  and  $L$ , then*

$$\delta(m) = \frac{\sqrt{1+m^2}}{k_1 \max\{1, |m|\} + k_2 \min\{1, |m|\}}. \quad (2)$$

*If  $K$  and  $L$  lie on a vertical line, then by definition,  $d_E(K, L) = \delta(m) d_G(K, L)$ .*

*Proof.* For any two points  $K = (x_1, y_1)$  and  $L = (x_2, y_2)$  with  $x_1 \neq x_2$ ; then  $m = (y_2 - y_1) / (x_2 - x_1)$ . Equation (2) is derived by a straightforward calculation with  $m$  and the coordinate definitions of  $d_E(K, L)$  and  $d_G(K, L)$  given in section 1.  $\square$

**Theorem 1.** *For any two distinct points  $K_1 = (x_1, y_1)$  and  $K_2 = (x_2, y_2)$  in the analytical plane, if  $L = (x, y)$  is a point on the line passing through  $K_1$  and  $K_2$ , then*

$$d_G[K_1L]/d_G[LK_2] = d_E[K_1L]/d_E[LK_2].$$

*That is, the ratios of the Euclidean and GAM directed lengths are the same.*

*Proof.* If  $L = K_1$  then both ratios are equal to 0. If  $L$  is at infinity then both ratios are equal to  $-1$ . Therefore without loss of generality, let  $K_1 \neq L \neq K_2$ . Then it's enough to show  $d_G(K_1, L) / d_G(L, K_2) = d_E(K_1, L) / d_E(L, K_2)$ . It's clear by Lemma 1. This completes proof.  $\square$

The following corollary shows how one can find the coordinates of the division point which divides the line segment joining two given points in a given ratio.

**Corollary 1.** For any two distinct points  $K_1 = (x_1, y_1)$  and  $K_2 = (x_2, y_2)$  in the GAM plane, if  $L = (x, y)$  divides the line segment  $K_1K_2$ , in ratio  $\delta$  then,

$$x = \frac{x_1 + \delta x_2}{1 + \delta}, \quad y = \frac{y_1 + \delta y_2}{1 + \delta}; \quad \delta \in \mathbb{R}, \quad \delta \neq -1$$

as in the Euclidean plane.

*Proof.* We would rather give a direct proof even though the Corollary follows from Theorem 1. The given formula is obvious when  $\delta = 0$  or  $\delta = -1$ . If  $\delta \neq 0, -1$  and  $L$  divides the line segment  $K_1K_2$  in ratio  $\delta$ , we have  $|d_G[K_1L_1]/d_G[L_1K_2]| = |\delta|$ . That is

$$\frac{k_1 \max\{|x_1 - x|, |y_1 - y|\} + k_2 \max\{|x_1 - x|, |y_1 - y|\}}{k_1 \max\{|x - x_2|, |y - y_2|\} + k_2 \max\{|x - x_2|, |y - y_2|\}} = |\delta|. \quad (3)$$

Since  $K_1 \neq K_2$ ,

Case (i) If  $|x_1 - x| \geq |y_1 - y|$  then using the definition of  $d_G$  distance we get

$$|\delta| = |\delta| \left( \frac{k_1 |x_1 - x_2| + k_2 |y_1 - y_2|}{k_1 |x_1 - x_2| + k_2 |y_1 - y_2|} \right) = \frac{k_1 |\delta x_1 - \delta x_2| + k_2 |\delta y_1 - \delta y_2|}{k_1 |x_1 - x_2| + k_2 |y_1 - y_2|}.$$

Adding  $x_1 - x_1$  and  $y_1 - y_1$  to the first and second summands in the numerator and similarly  $\delta x_2 - \delta x_2$  and  $\delta y_2 - \delta y_2$  in the denominator respectively, one obtains

$$|\delta| = \frac{k_1 |\delta x_1 + x_1 - x_1 - \delta x_2| + k_2 |\delta y_1 + y_1 - y_1 - \delta y_2|}{k_1 |x_1 + \delta x_2 - \delta x_2 - x_2| + k_2 |y_1 + \delta y_2 - \delta y_2 - y_2|}.$$

Multiplying the numerator and the denominator of the right side of the last statement by  $1/|1 + \delta|$ , one gets

$$\begin{aligned} |\delta| &= \frac{k_1 \left| \frac{\delta x_1 + x_1 - x_1 - \delta x_2}{1 + \delta} \right| + k_2 \left| \frac{\delta y_1 + y_1 - y_1 - \delta y_2}{1 + \delta} \right|}{k_1 \left| \frac{x_1 + \delta x_2 - \delta x_2 - x_2}{1 + \delta} \right| + k_2 \left| \frac{y_1 + \delta y_2 - \delta y_2 - y_2}{1 + \delta} \right|} \\ &= \frac{k_1 \left| \frac{(1 + \delta)x_1}{1 + \delta} - \frac{x_1 + \delta x_2}{1 + \delta} \right| + k_2 \left| \frac{(1 + \delta)y_1}{1 + \delta} - \frac{y_1 + \delta y_2}{1 + \delta} \right|}{k_1 \left| \frac{x_1 + \delta x_2}{1 + \delta} - \frac{(1 + \delta)x_2}{1 + \delta} \right| + k_2 \left| \frac{y_1 + \delta y_2}{1 + \delta} - \frac{(1 + \delta)y_2}{1 + \delta} \right|} \\ &= \frac{k_1 \left| x_1 - \frac{x_1 + \delta x_2}{1 + \delta} \right| + k_2 \left| y_1 - \frac{y_1 + \delta y_2}{1 + \delta} \right|}{k_1 \left| \frac{x_1 + \delta x_2}{1 + \delta} - x_2 \right| + k_2 \left| \frac{y_1 + \delta y_2}{1 + \delta} - y_2 \right|} = \frac{k_1 |x_1 - x| + k_2 |y_1 - y|}{k_1 |x_2 - x| + k_2 |y_2 - y|}. \end{aligned}$$

Comparing this result with equation (2) we obtain

$$x = \frac{x_1 + \delta x_2}{1 + \delta}, \quad y = \frac{y_1 + \delta y_2}{1 + \delta}.$$

Case (ii) If  $|y_1 - y| < |x_1 - x|$  then it can be proved similarly case (i).  $\square$

Figure 1:

### 3 Theorems of Menelaus and Ceva in the GAM Plane

In this section, we study the GAM version of the Theorems of Menelaus and Ceva. Indeed, the validity of these theorems is obvious from the Theorem 1, but we prefer to state and give partial proofs for them.

**Theorem 2.** (*Menelaus'es Theorem.*) *Let  $\{K_1, K_2, K_3\}$  be a triangle and  $L_1, L_2, L_3$  be on the lines that contain the sides  $K_1K_2, K_2K_3, K_3K_1$  respectively, in the GAM plane. If  $L_1, L_2, L_3$  are collinear, then*

$$\frac{d_G[K_1L_1]}{d_G[L_1K_2]} \cdot \frac{d_G[K_2L_2]}{d_G[L_2K_3]} \cdot \frac{d_G[K_3L_3]}{d_G[L_3K_1]} = -1$$

where none of  $L_1, L_2, L_3$  coincide with any of  $K_1, K_2, K_3$ .

*Proof.* According to the positions of points  $K_1, K_2, K_3$  and  $L_1, L_2, L_3$ , there are several possible cases. We give a proof of the theorem only in the following special case.

Let  $K_i = (x_i, y_i)$ ,  $i = 1, 2, 3$  and  $x_i \neq y_{i+1}$  and let  $L_1, L_2, L_3$  be on a line  $l$  given by  $y = mx + k$  such that  $L_i = l \wedge K_iK_{i+1(mod3)}$  and  $l$  is not parallel to the line  $K_iK_{i+1}$ , for  $i = 1, 2, 3$  (see Figure 1). Clearly  $mx_i - y_i + k \neq 0$  since  $K_i \neq L_j$  for  $i, j = 1, 2, 3$  and  $m \neq (y_{i+1} - y_i)(x_{i+1} - x_i)^{-1}$ . The equation of the line  $K_iK_{i+1}$  is given by

$$y = \frac{(y_{i+1} - y_i)}{(x_{i+1} - x_i)}x - \frac{(x_iy_{i+1} - x_{i+1}y_i)}{(x_{i+1} - x_i)}.$$

It follows from a simple calculation that

$$\begin{aligned} L_i &= \left( \frac{x_iy_{i+1} - x_{i+1}y_i - kx_i + kx_{i+1}}{mx_i - mx_{i+1} - y_i + y_{i+1}}, \frac{mx_iy_{i+1} - mx_{i+1}y_i - ky_i + ky_{i+1}}{mx_i - mx_{i+1} - y_i + y_{i+1}} \right) \\ &= (u, v). \end{aligned}$$

Now let us find  $\frac{d_G[K_iL_i]}{d_G[L_iK_{i+1}]} \cdot \frac{d_G[K_1L_1]}{d_G[L_1K_2]} = -\frac{d_G(K_1, L_1)}{d_G(L_1, K_2)}$ . Since  $K_1, K_2$  and  $L_1$  are collinear,

Case (i) If  $|x_1 - u| \geq |y_1 - v|$  then using the definition of  $d_G$ - distance one gets

$$\begin{aligned}
&= \frac{k_1 \left| x_1 - \frac{x_1 y_2 - x_2 y_1 - k x_1 + k x_2}{m x_1 - m x_2 - y_1 + y_2} \right| + k_2 \left| y_1 - \frac{m x_1 y_2 - m x_2 y_1 - k y_1 + k y_2}{m x_1 - m x_2 - y_1 + y_2} \right|}{k_1 \left| \frac{x_1 y_2 - x_2 y_1 - k x_1 + k x_2}{m x_1 - m x_2 - y_1 + y_2} - x_2 \right| + k_2 \left| y_1 - \frac{m x_1 y_2 - m x_2 y_1 - k y_1 + k y_2}{m x_1 - m x_2 - y_1 + y_2} - y_2 \right|} \\
&= \frac{k_1 |x_1 (m x_1 - y_1 + k) - x_2 (m x_1 - y_1 + k)| + k_2 |y_1 (m x_1 - y_1 + k) - y_2 (m x_1 - y_1 + k)|}{k_1 |x_2 (m x_2 - y_2 + k) - x_1 (m x_2 - y_2 + k)| + k_2 |y_2 (m x_2 - y_2 + k) - y_1 (m x_2 - y_2 + k)|} \\
&= \frac{|m x_1 - y_1 + k| (k_1 |x_1 - x_2| + k_2 |y_1 - y_2|)}{|m x_2 - y_2 + k| (k_1 |x_2 - x_1| + k_2 |y_2 - y_1|)} \\
&= \frac{|m x_1 - y_1 + k|}{|m x_2 - y_2 + k|}.
\end{aligned}$$

Similarly,

$$\frac{d_G[K_2 L_2]}{d_G[L_2 K_3]} = \frac{|m x_2 - y_2 + k|}{|m x_3 - y_3 + k|}$$

and

$$\frac{d_G[K_3 L_3]}{d_G[L_3 P K_1]} = \frac{|m x_3 - y_3 + k|}{|m x_1 - y_1 + k|}$$

and consequently,

$$\frac{d_G[K_i L_i]}{d_G[L_i K_{i+1}]} = s \frac{|m x_i - y_i + k|}{|m x_{i+1} - y_{i+1} + k|}, \quad s = \begin{cases} -1, & \text{if } i = 1 \\ 1, & \text{if } i = 2, 3. \end{cases}$$

Now , it can be easily computed that

$$\prod_{i=1}^3 (d_G[K_i L_i] / d_G[L_i K_{i+1}]) = -1.$$

Case (ii) If  $|x_1 - u| < |y_1 - v|$  then it can be proved similarly case (i).  $\square$

**Theorem 3.** (Converse of Menelaus'es Theorem.) Let  $\{K_1, K_2, K_3\}$  be a triangle and  $L_1, L_2, L_3$  be three points on the lines that contain the sides  $K_1 K_2, K_2 K_3, K_3 K_1$  respectively, in the GAM plane. If

$$\frac{d_G[K_1 L_1]}{d_G[L_1 K_2]} \cdot \frac{d_G[K_2 L_2]}{d_G[L_2 K_3]} \cdot \frac{d_G[K_3 L_3]}{d_G[L_3 K_1]} = -1,$$

then  $L_1, L_2, L_3$  are collinear. Note that none of  $L_1, L_2, L_3$  coincide with any of  $K_1, K_2, K_3$ .

**Theorem 4.** (Ceva's Theorem.) Let  $\{K_1, K_2, K_3\}$  be a triangle and lines  $l_1, l_2, l_3$  pass through the vertices  $K_1, K_2, K_3$ , respectively and intersect lines containing the opposite sides at points  $L_1, L_2, L_3$ . The lines  $l_1, l_2, l_3$  are concurrent (or parallel) if and only if

$$\frac{d_G[K_1 L_3]}{d_G[L_3 K_2]} \cdot \frac{d_G[K_2 L_1]}{d_G[L_1 K_3]} \cdot \frac{d_G[K_3 L_2]}{d_G[L_2 K_1]} = 1.$$

Note that none of  $L_1, L_2, L_3$  are  $K_1, K_2, K_3$ .

Figure 2:

## 4 Theorems of Directed Lines(Strahlensätze).

In general, the axiom congruence and consequently properties of similarity for triangles are not valid in the GAM plane . But, it follows from Theorem 1 that the following *directed line theorem* [18] is valid in it.

**Theorem 5.** *Let a pencil of lines be intersected by a family of parallel lines in the GAM plane (see Figure 2)*

(i) *The ratios of directed lengths of the corresponding segments on the lines belonging to the pencil are the same. For example*

$$\begin{aligned} d_G[SP] : d_G[SQ] : d_G[SR] &= d_G[SP_1] : d_G[SQ_1] : d_G[SR_1] \\ &= d_G[SP_2] : d_G[SQ_2] : d_G[SR_2] \end{aligned}$$

or

$$d_G[SP_1] : d_G[SQ_1] = d_G[P_1P_2] : d_G[Q_1Q_2].$$

(ii) *The ratios of directed lengths of line segments on the parallel lines and corresponding segments on the lines belonging to the pencil, which are measured from the vertex, are the same. For example*

$$\begin{aligned} d_G[RQ] : d_G[R_1Q_1] : d_G[R_2Q_2] &= d_G[SR] : d_G[SR_1] : d_G[SR_2] \\ &= d_G[SQ] : d_G[SQ_1] : d_G[SQ_2] \end{aligned}$$

or

$$\begin{aligned} d_g[PQ] : d_g[P_1Q_1] : d_g[P_2Q_2] &= d_g[SP] : d_g[SP_1] : d_g[SP_2] \\ &= d_g[SQ] : d_g[SQ_1] : d_g[SQ_2]. \end{aligned}$$

(iii) *The ratios of directed lengths of the corresponding segments on the parallel lines are the same. That is*

$$d_G[PQ] : d_G[QR] = d_G[P_1Q_1] : d_G[Q_1R_1] = d_G[P_2Q_2] : d_G[Q_2R_2].$$

Notice that here  $p : q : r = p_1 : q_1 : r_1$  if and only if  $p/p_1 = q/q_1 = r/r_1$ .

**Acknowledgements.** We thank Professor Rüstem Kaya and Professor Münevver Özcan for providing us with some references.

## References

- [1] Z. Akça, A. Bayar, S. Ekmekçi, *The Norm in CC-Plane Geomtry*, Pi Mu Epsilon Journal, 12(6), (2007), 321-324
- [2] Z. Akça, R. Kaya, *On the Taxicab Trigonometry*, Jour. of Inst. of Math.Comp. Sci.(Math. Ser.), 10, (1997), 151-159.
- [3] A. Bayar, S. Ekmekçi, Z. Akça, *On the Plane Geometry with Generalized Absolute Value Metric*, Mathematical Problems in Engineering, Vol. 2008, 1-8, DOI: 10.1155/2008/673275.
- [4] A. Bayar, S. Ekmekçi, *On The Chinese Checker Sine and Cosine Functions*, International Journal of Mathematics and Analysis, 1(3), (2006), 249-254.
- [5] H.B. Çolakođlu, R. Kaya, *Taxicab Versions of The Pythagorean Theorem*, Pi Mu Epsilon Journal, 12(9), (2008), No. 9, 535-539.
- [6] H.B. Çolakođlu, R. Kaya, *On The Regular Polygons in the Chinese Checker Plane*, Applied Sciences, 10,(2008), 29-37.
- [7] H.B. Çolakođlu, R. Kaya, *A Synthetic Approach to the Taxicab Circles*, Applied Science, 9, (2007), 67-77.
- [8] B. Dijvak, *Notes on Taxicab Geometry*, Scientific and Professional Information Journal Of Crotian Society for Constructive Geometry and Computer Graphics, 5, (2000), 5-9.
- [9] C. Ekici, I. Kocayusufođlu, Z. Akça, *The Norm in Taxicab Geometry*, Turkish Journal of Mathematics, 22, (1998), 295-307.
- [10] Ö. Gelişgen, R. Kaya, *CC-Analog of the Theorem of Pythagorean*, Algebras Groups Geom., 23(2), (2006), 179-188.
- [11] Ö. Gelişgen, R. Kaya, M. Özcan, *Distance Formulae in the Chinese Checker Space*, Int. Jour. Of Pure and Appl. Math., (26)1, (2006), 35-44.
- [12] Y.P. Ho, Y. Liu, *Parabolas in Taxicab Geometry*, Missouri J. of Math. Sci., 8, (1996), 63-72.
- [13] R. Kaya, Ö. Gelişgen, S. Ekmekçi, A. Bayar, *Isometries of the plane with generalized Absolute Value Metric*, Rocky Mountain Journal of Mathematics, 39(2), (2009), 573-591.
- [14] R. Kaya, *Area Formula for Taxicab Triangles*, Pi Mu Epsilon Journal, 12, (2006), 213-220.
- [15] R. Kaya, H.B. Çolakođlu, *Taxicab Versions of Some Euclidean Theorems*, Int. Jour. Of Pure and Appl. Math., 26(1), (2006), 69-81.



- [16] R. Kaya, Ö. Gelisgen, S. Ekmekçi, A. Bayar, *On the Group Isometries of CC-Plane*, Missouri J. Math. Sci., 3, (2006), 221-233.
- [17] E.F. Krause, *Taxicab Geometry*, Addison-Wesley, Menlo Park, California, Dover Publications, New York, 1987.
- [18] H. Kreul, K. Kulke, H. Pester, R. Schroedter, *Mathematik Leicht Gemacht*, Verlag Harri Deutsch-Thun-Frankfurt am Main, 1990.
- [19] R. Laatsch, *Prymidal Sections in Taxicab Geometry*, Mathematics Magazine, 55, (1982), 205-212.
- [20] K. Menger, *You Will Like Geometry*, Guildbook of the Illinois Institute of Technology Geometry Exhibit, Museum of Science and Industry, Chicago, IL, 1952.
- [21] R.S. Milmann, G.D. Parker, *Geometry; A metric approach with models*, Springer, 1991.
- [22] H. Minkowski, *Gesammelte Abhandlungen*, Chelsea Publishing Co., New York, 1967.
- [23] M. Özcan, S. Ekmekçi, A. Bayar, *A note on the Variation of the Taxicab Lengths Under Rotations*, Pi Mu Epsilon Journal, 11(7), (2002), 381-384.
- [24] M. Özcan, R. Kaya, *Area of a Triangle in terms of the Taxicab Distance*, Missouri J. Math. Sci., 15, (2003), 175-185.
- [25] M. Özcan, R. Kaya, *On the Ratio of Directed Lengths in the Taxicab Plane and Related Properties*, Missouri J. Math. Sci., 14, (2002), 107-117.
- [26] D.J. Schattschneider, *The Taxicab Group*, Amer Math. Monthly., 91, (1984), 423-428.
- [27] M. Turan, M. Özcan, *Two-Foci Chinese Checker Hyperbolas*, Int. Jour. Of Pure and Appl. Math., 16(4), (2004), 509-520.
- [28] M. Turan, M. Özcan, *Two-Foci Chinese Checker Ellipses*, Int. Jour. Of Pure and Appl. Math., 16(1), (2004), 119-127.
- [29] M. Turan, M. Özcan, *General Equation for Chinese Checker Conics and Focus-Directrix Chinese Checker Conics*, Int. Jour. Of Pure and Appl. Math., 30(3), (2006), 397-406.

Author 1: Gökhan Soydan  
Işıklar Air Force High School 16039, Bursa, TURKEY  
E-mail: gsoydan@uludag.edu.tr

Author 2: Yusuf Dođru  
Işıklar Air Force High School 16039, Bursa, TURKEY

*E-mail:* yusufdogru1@yahoo.com

Author 3: N.Umut Arslandođan  
Işıklar Air Force High School 16039, Bursa, TURKEY  
*E-mail:* nuarslandogan@gmail.com