

## Polynomials of the Laguerre type

Gospava B. Djordjević<sup>a</sup>

<sup>a</sup>University of Niš, Faculty of Technology, 16000 Leskovac, Serbia

**Abstract.** In this note we shall study a class of polynomials  $\{f_{n,m}^{c,r}(x)\}$ , where  $c$  is some real number,  $r \in \mathbb{N} \cup \{0\}$ ,  $m \in \mathbb{N}$ . These polynomials are defined by the generating function. Also, for these polynomials we find an explicit representation in the form of the hypergeometric function; some identities of the convolution type are presented; some special cases are shown. The special cases of these polynomials are: Panda's polynomials [2], [4]; the generalized Laguerre polynomials [1], [6]; the Celine Fasenmyer polynomials [3].

### 1. Polynomials $f_{n,m}^{c,r}(x)$

Let  $\phi(u)$  be a formal power-series expansion

$$\phi(u) = \sum_{n=0}^{\infty} \gamma_n u^n, \quad \gamma_n = \frac{(-r^r)^n}{n!}. \quad (1.1)$$

We define the polynomials  $f_{n,m}^{c,r}(x)$  as

$$(1 - t^m)^{-c} \phi\left(\frac{-4xt}{(1 - t^m)r}\right) = \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x) t^n. \quad (1.2)$$

We prove the following result.

**Theorem 1.1.** *The polynomials  $f_{n,m}^{c,r}(x)$  satisfy the following relations:*

$$f_{n,m}^{c,r}(x) = \frac{(4r^r x)^n}{n!} {}_{(r+1)m}F_{rm-1} \times A, \quad (1.3)$$

where

$$A = \left[ \begin{array}{c} -\frac{n}{m}, \dots, \frac{m-1-n}{m}, \frac{1-c-rn}{rm}, \dots, \frac{rm-c-rn}{rm}, \frac{(-1)^{m-1}(4r^r x)^{-m}(rm)^m}{(rm-1)^{m-1}} \\ \frac{1-c-rn}{rm-1}, \frac{2-c-rn}{rm-1}, \dots, \frac{rm-1-c-rn}{rm-1} \end{array} \right] \quad (1.4)$$

$$x^n = \frac{n!}{4^n (r^r)^n} \sum_{k=0}^{\lfloor n/m \rfloor} (-1)^k \binom{c + rn - rmk}{k} f_{n-mk,m}^{c,r}(x). \quad (1.5)$$

2010 *Mathematics Subject Classification.* Primary 11B83; Secondary 11B37, 11B39.

*Keywords.* Generating function; Hypergeometric function; Convolution.

Received: 25 June 2011; Accepted: 12 August 2011

Communicated by Dragan S. Djordjević

Research supported by Ministry Education and Science, Republic of Serbia, Grant No. 174007

*Email address:* [gospava48@ptt.rs](mailto:gospava48@ptt.rs) (Gospava B. Djordjević)

Proof. Using (1.1) and (1.2), we find:

$$\begin{aligned} (1 - t^m)^{-c} \phi\left(\frac{-4xt}{(1 - t^m)^r}\right) &= (1 - t^m)^{-c} \sum_{n=0}^{\infty} \frac{(-r^r)^n}{n!} \frac{(-4x)^n t^n}{(1 - t^m)^{rn}} = \sum_{n=0}^{\infty} \frac{(4r^r x)^n t^n}{n!} (1 - t^m)^{-c-rn} \\ &= \left(\sum_{n=0}^{\infty} \frac{(4r^r x)^n t^n}{n!}\right) \left(\sum_{k=0}^{\infty} \binom{-c-rn}{k} (-t^m)^k\right) = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{(4r^r x)^{n-k} t^{n-k}}{(n-k)!} \binom{-c-r(n-k)}{k} (-1)^k t^{mk} \\ &\quad (n - k - mk := n, \quad n - k := n - mk, \quad k \leq [n/m]) \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} \frac{(-1)^k (4r^r)^{n-mk} x^{n-mk}}{(n-mk)!} \binom{-c-r(n-mk)}{k} t^n = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{[n/m]} \frac{(4r^r)^{n-mk} x^{n-mk} (c+r(n-mk))_k}{k!(n-mk)!}\right) t^n. \end{aligned}$$

Using the well-known equalities ([5])

$$(\alpha)_{n+k} = (\alpha)_n (\alpha + n)_k, \quad \frac{(-1)^k}{(n-k)!} = \frac{(-n)_k}{n!}, \quad (\alpha)_{n-k} = \frac{(-1)^k (\alpha)_n}{(1 - \alpha - n)_k},$$

we find that

$$\begin{aligned} \frac{(c+r(n-mk))_k}{(n-mk)!} &= \frac{(c)_{r(n-mk)+k}}{(c)_{r(n-mk)}} \cdot \frac{(-1)^{mk} (-1)^{mk}}{(n-mk)!} = \frac{(-1)^{(rm-1)k} (c)_{rn}}{(1-c-rn)_{(rm-1)k}} \cdot \frac{(1-c-rn)_{rmk}}{(-1)^{rmk} (c)_{rn}} \cdot \frac{(-1)^{mk} (-n)_{mk}}{n!} \\ &= \frac{(-1)^{(m-1)k} (1-c-rn)_{rmk} (-n)_{mk}}{(1-c-rn)_{(rm-1)k} n!} = \frac{(-1)^{(m-1)k} (rm)^{rmk} m^{mk} \cdot A \cdot B}{(rm-1)^{(rm-1)k} n! \cdot C} \end{aligned}$$

where

$$\begin{aligned} A &= \left(\frac{1-c-rn}{rm}\right)_k \cdot \left(\frac{2-c-rn}{rm}\right)_k \cdots \left(\frac{rm-c-rn}{rm}\right)_k, \\ B &= \left(\frac{-n}{m}\right)_k \cdot \left(\frac{1-n}{m}\right)_k \cdots \left(\frac{m-1-n}{m}\right)_k, \\ C &= \left(\frac{1-c-rn}{rm-1}\right)_k \cdot \left(\frac{2-c-rn}{rm-1}\right)_k \cdots \left(\frac{rm-1-c-rn}{rm-1}\right)_k. \end{aligned}$$

From the other side, because of the next equality,

$$(1 - t^m)^{-c} \phi\left(\frac{-4xt}{(1 - t^m)^r}\right) = \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x) t^n$$

it follows

$$f_{n,m}^{c,r}(x) = \frac{(4r^r x)^n}{n!} {}_{(r+1)m}F_{rm-1} \left[ \begin{matrix} 1-c-rn, \dots, \frac{rm-c-rn}{rm}, \frac{-n}{m}, \dots, \frac{m-1-n}{m}, \frac{(-1)^{m-1} (4r^r x)^{-m} (rm)^{rm}}{(rm-1)^{rm-1}} \\ \frac{1-c-rn}{rm-1}, \frac{2-c-rn}{rm-1}, \dots, \frac{rm-1-c-rn}{rm-1} \end{matrix} \right].$$

These are the required equalities (1.3) and (1.4).

Again, from (1.1) and (1.2), we get:

$$\begin{aligned} \sum_{n=0}^{\infty} (-4x)^n t^n \frac{(-r^r)^n}{n!} &= (1 - t^m)^{c+rn} \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x) t^n = \left(\sum_{k=0}^{\infty} \binom{c+rn}{k} (-1)^k t^{mk}\right) \left(\sum_{n=0}^{\infty} f_{n,m}^{c,r}(x) t^n\right) \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} (-1)^k \binom{c+r(n-mk)}{k} f_{n-mk,m}^{c,r}(x) t^n. \end{aligned}$$

Hence, we get

$$x^n = \frac{n!}{4^n (r^r)^n} \sum_{k=0}^{[n/m]} (-1)^k \binom{c+r(n-mk)}{k} f_{n-mk,m}^{c,r}(x),$$

which yields the equality (1.5).  $\square$

## 2. Some special cases of $f_{n,m}^{c,r}(x)$

If  $r = 1$  and  $m > 1$ , then (1.3)–(1.4) become

$$f_{n,m}^{c,1}(x) = \frac{(-4x)^n}{n!} {}_2F_{m-1} \left[ \begin{matrix} \frac{1-c-n}{m}, \frac{2-c-n}{m}, \dots, \frac{m-c-n}{m}, \frac{-n}{m}, \frac{1-n}{m}, \dots, \frac{m-1-n}{m}, \frac{(\frac{m}{4x})^m}{(1-m)^{m-1}} \\ \frac{1-c-n}{m-1}, \frac{2-c-n}{m-1}, \dots, \frac{m-1-c-n}{m-1} \end{matrix} \right].$$

If  $m = 1$  and  $r > 1$ , then (1.3)–(1.4) yield

$$f_{n,1}^{c,r}(x) = \frac{(4r^r x)^n}{n!} {}_{r+1}F_{r-1} \left[ \begin{matrix} \frac{1-c-rn}{r}, \dots, \frac{r-c-rn}{r}, -n, \frac{(4r^r x)^{-1}}{(r-1)^{r-1}} \\ \frac{1-c-rn}{r-1}, \dots, \frac{r-1-c-rn}{r-1} \end{matrix} \right].$$

For  $r = 0$  in (1.2), and  $\phi(u) = e^u$ , we have  $(1 - t^m)^{-c} \phi(-4xt) = \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x) t^n$ , and hence we get the following equalities:

$$e^{-4xt} = (1 - t^m)^c \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x) t^n,$$

and also

$$\sum_{n=0}^{\infty} \frac{(-4x)^n t^n}{n!} = \left( \sum_{k=0}^{\infty} \binom{c}{k} (-t^m)^k \right) \left( \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x) t^n \right) = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} (-1)^k \binom{c}{k} f_{n-mk,m}^{c,0}(x) t^n.$$

So, we get

$$x^n = \frac{n!}{(-4)^n} \sum_{k=0}^{[n/m]} (-1)^k \binom{c}{k} f_{n-mk,m}^{c,0}(x). \tag{2.1}$$

For  $c = 0$  in (1.3) and (1.4), we get the following formula

$$f_{n,m}^{0,r}(x) = \frac{(4r^r x)^n}{n!} {}_{(r+1)m}F_{rm-1} \left[ \begin{matrix} \frac{1-rn}{rm}, \dots, \frac{rm-rn}{rm}, \frac{-n}{m}, \dots, \frac{m-1-n}{m}, \frac{(-1)^{m-1} (4r^r x)^{-m} m^m}{(rm-1)^{m-1}} \\ \frac{1-rn}{rm-1}, \dots, \frac{rm-1-rn}{rm-1} \end{matrix} \right].$$

For  $c = 1$ ,  $m = 2$  and  $r = 2$ , then,  $\gamma_n = \frac{(-4)^n}{n!}$  and by (1.3) and (1.4), we get the following formula

$$f_{n,2}^{1,2}(x) = \frac{(4x)^n}{n!} {}_6F_3 \left[ \begin{matrix} \frac{-2n}{4}, \frac{1-2n}{4}, \frac{2-2n}{4}, \frac{3-2n}{4}, \frac{-n}{2}, \frac{1-n}{2}, \frac{-1}{12^3 x^2} \\ \frac{-2n}{3}, \frac{1-2n}{3}, \frac{2-2n}{3} \end{matrix} \right]$$

Note that the generalized Laguerre polynomials are the special case of the polynomials  $f_{n,m}^{c,r}(x)$ , that is,  $L_{n,m}^c(x) = f_{n,m}^{c-1,1}(x/4)$ . So, we get the following representation

$$L_{n,m}^c = \frac{x^n}{n!} {}_2mF_{m-1} \left[ \begin{matrix} \frac{2-c-n}{m}, \dots, \frac{m+1-c-n}{m}, \frac{-n}{m}, \dots, \frac{m-1-n}{m}, \frac{x^{-m}m^m}{(1-m)^{m-1}} \\ \frac{2-c-n}{m-1}, \dots, \frac{m-c-n}{m-1} \end{matrix} \right].$$

For the Laguerre polynomials  $L_{n,1}^c(x) \equiv L_n^c(x)$ , where

$$(1-t)^{-c} \exp\left\{\frac{-xt}{1-t}\right\} = \sum_{n=0}^{\infty} L_n^c(x) \frac{t^n}{n!},$$

the following statement holds.

**Theorem 2.1.** Let  $D = \frac{d}{dx}$  and  $g \in C^\infty(-\infty, +\infty)$  and  $g(x) \neq 0$ , then

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} D^{n-k-j}\{g^{-1}\} D^j\{g\} = (-1)^n n!. \tag{2.2}$$

*Proof.* Using the known formula ([5])

$$DL_n^c(x) = DL_{n-1}^c(x) - L_{n-1}^c(x), \quad n \geq 1,$$

we get the following equalities:

$$DL_n^c(x) = (D-1)L_{n-1}^c(x), \quad D^2L_n^c(x) = (D-1)^2L_{n-2}^c(x), \quad \dots, \quad D^sL_n^c(x) = (D-1)^sL_{n-s}^c(x), \quad n \geq s.$$

Hence, for  $s = n$ , we obtain that

$$D^n L_n^c(x) = \left( \sum_{k=0}^n (-1)^k \binom{n}{k} D^{n-k} \right) \{1\} = \sum_{k=0}^n (-1)^k \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} D^{n-k-j}\{g^{-1}\} D^j\{g\}.$$

Since  $D^n L_n^c(x) = (-1)^n n!$ , we get

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} D^{n-k-j}\{g^{-1}\} D^j\{g\} = (-1)^n n!,$$

which leads to (2.2).  $\square$

Depending on the chosen functions  $g(x)$  and from (2.2), we get some interesting relations.

1° For  $g(x) = e^{ax}$ ,  $a$  is any rial number, and  $g^{-1}(x) = e^{-ax}$ , we get

$$a^n \sum_{k=0}^n \frac{a^{-k}}{k!} \sum_{j=0}^{n-k} \frac{(-1)^j}{j!(n-k-j)!} = 1.$$

2° If  $g(x) = (1+x)^\alpha$ , for  $x > -1$ ,  $\alpha \neq 0$ , then we get

$$(\alpha)_n \Gamma(\alpha) \sum_{k=0}^n \frac{(1+x)^k}{k!} \sum_{j=0}^{n-k} \frac{1}{(1-\alpha-n)_{k+j-2} \Gamma(\alpha-j+2)} = \frac{(1+x)^n}{\alpha n!},$$

or

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \frac{(-1)^j (1+x)^k \left( \prod_{i=0}^j (\alpha+1-i) \right) \left( \prod_{s=0}^{n-k-j-1} (\alpha+s) \right)}{k! j! (n-k-j)!} = \frac{(1+x)^n}{\alpha}.$$

3° For  $g(x) = a^x$ ,  $a > 0$  and  $a \neq 1$ , we obtain

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \frac{(-1)^j (\ln a)^{n-k}}{j! k! (n-k-j)!} = 1.$$

4° For  $g(x) = x^\alpha e^x$ , we have the following formula

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i=0}^{n-k-j} \sum_{l=0}^j \frac{(-1)^j (\alpha)_j (\alpha+1-l)_l x^{-i-l}}{k! i! l! (n-k-j)! (j-l)!} = 1,$$

or in the form

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i=0}^{n-k-j} \sum_{l=0}^j \frac{(-1)^{k+i} (-n)_{k+j+i} (\alpha)_j}{x^{i+l} k! i! l! (j-l)!} = \frac{n!}{\Gamma(\alpha+1)}.$$

Hence, for  $\alpha = n$ ,  $n \in \mathbb{N}$ , we get

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i=0}^{n-k-j} \sum_{l=0}^j \frac{(-1)^{k+i} (-n)_{k+j+i} j!}{x^{i+l} i! l! (j-l)!} = 1.$$

5° For  $g(x) = x^\alpha$ ,  $x \geq 0$  and  $x \neq 1$ ,  $\alpha \neq 0$ , we obtain

$$\sum_{k=0}^n \sum_{j=0}^{n-k} \frac{(-1)^j \prod_{i=0}^{n-k-j-1} (\alpha+j) \prod_{s=0}^{j-1} (\alpha-s) x^k}{k! j!} = x^n.$$

### 3. Some identities of the convolution type

Using the following equality

$$(1 - t^m)^{-c/2} \phi\left(\frac{-4xt}{(1 - t^m)^r}\right) = (1 - t^m)^{c/2} \sum_{n=0}^{\infty} f^{c/2,r}(x)t^n, \quad (3.1)$$

and by (1.1) and (1.2), we get

$$\sum_{n=0}^{\infty} f_{n,m}^{c/2,r}(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{c/2}{k} (-1)^k f_{n-k,m}^{c/2,r}(x)t^{n+mk-k} = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} \binom{c/2}{k} (-1)^k f_{n-mk,m}^{c/2,r}(x)t^n.$$

Hence, we get

$$f_{n,m}^{c/2,r}(x) = \sum_{k=0}^{[n/m]} (-1)^k \binom{c/2}{k} f_{n-mk,m}^{c/2,r}(x). \quad (3.2)$$

Again, by (1.1) and (1.2), for  $\phi(u) = e^u$  we find:

$$(1 - t^m)^{-c} \phi\left(\frac{-4(x_1 + x_2 + \dots + x_s)t}{(1 - t^m)^r}\right) = \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x_1 + x_2 + \dots + x_s)t^n,$$

that is,

$$(1 - t^m)^{-c/s} e^{\frac{-4x_1 t}{(1-t^m)^r}} \dots (1 - t^m)^{-c/s} e^{\frac{-4x_s t}{(1-t^m)^r}} = \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x_1 + \dots + x_s)t^n,$$

hence

$$\left(\sum_{n=0}^{\infty} f_{n,m}^{c/s,r}(x_1)t^n\right) \left(\sum_{n=0}^{\infty} f_{n,m}^{c/s,r}(x_2)t^n\right) \dots \left(\sum_{n=0}^{\infty} f_{n,m}^{c/s,r}(x_s)t^n\right) = \sum_{n=0}^{\infty} f_{n,m}^{c,r}(x_1 + \dots + x_s)t^n.$$

So, we get

$$\sum_{i_1 + \dots + i_s = n} f_{i_1,m}^{c/s,r}(x_1) f_{i_2,m}^{c/s,r}(x_2) \dots f_{i_s,m}^{c/s,r}(x_s) = f_{n,m}^{c,r}(x_1 + x_2 + \dots + x_s). \quad (3.3)$$

Let  $c = c_1 + \dots + c_k$  and  $x = x_1 + \dots + x_k$ , then we have, at, on the one side

$$(1 - t^m)^{-c_1 - \dots - c_k} \phi\left(\frac{-4(x_1 + \dots + x_k)t}{(1 - t^m)^r}\right) = \sum_{n=0}^{\infty} f_{n,m}^{c_1 + \dots + c_k,r}(x_1 + \dots + x_k)t^n,$$

and on the other hand

$$\begin{aligned} (1 - t^m)^{-c_1 - \dots - c_k} \phi\left(\frac{-4(x_1 + \dots + x_k)t}{(1 - t^m)^r}\right) &= \left(\sum_{n=0}^{\infty} f_{n,m}^{c_1,r}(x_1)t^n\right) \dots \left(\sum_{n=0}^{\infty} f_{n,m}^{c_k,r}(x_k)t^n\right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{i_1 + \dots + i_k = n} f_{i_1,m}^{c_1,r}(x_1) \dots f_{i_k,m}^{c_k,r}(x_k)\right) t^n. \end{aligned}$$

Hence we get the following formula

$$\sum_{i_1+\dots+i_k=n} f_{i_1,m}^{c_1,r}(x_1) \cdots f_{i_k,m}^{c_k,r}(x_k) = f_{n,m}^{c_1+\dots+c_k,r}(x_1+\dots+x_k). \quad (3.4)$$

If  $x_1 = x_2 = \dots = x_s = \frac{x}{s}$ , then (3.3) becomes

$$\sum_{i_1+\dots+i_s=n} f_{i_1,m}^{c/s,r}(x/s) \cdots f_{i_s,m}^{c/s,r}(x/s) = f_{n,m}^{c,r}(x), \quad (3.5)$$

where  $s$  is a natural number.

For  $r = 0$  in (1.2), and  $\phi(u) = e^u$ , we have  $(1 - t^m)^{-c} \phi(-4xt) = \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x)t^n$ , whence we get

$$e^{-4xt} = (1 - t^m)^c \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x)t^n,$$

and also

$$\sum_{n=0}^{\infty} \frac{(-4x)^n t^n}{n!} = \left( \sum_{k=0}^{\infty} \binom{c}{k} (-t^m)^k \right) \left( \sum_{n=0}^{\infty} f_{n,m}^{c,0}(x)t^n \right) = \sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor n/m \rfloor} (-1)^k \binom{c}{k} f_{n-mk,m}^{c,0}(x)t^n.$$

So, we get

$$x^n = \frac{n!}{(-4)^n} \sum_{k=0}^{\lfloor n/m \rfloor} (-1)^k \binom{c}{k} f_{n-mk,m}^{c,0}(x). \quad (3.6)$$

For  $c = 0$  in (1.3) and (1.4), we get the following formula

$$f_{n,m}^{0,r}(x) = \frac{(4r^r x)^n}{n!} {}_{(r+1)m}F_{rm-1} \left[ \begin{matrix} \frac{1-rn}{rm}, \dots, \frac{rm-rn}{rm}, \frac{-n}{m}, \dots, \frac{m-1-n}{m}, \frac{(-1)^{m-1}(4r^r x)^{-m} m^m}{(rm-1)^{m-1}} \\ \frac{1-rn}{rm-1}, \dots, \frac{rm-1-rn}{rm-1} \end{matrix} \right].$$

## References

- [1] G. B. Djordjević, *On the generalized Laguerre polynomials*, Fibonacci Quart. 39 (2001), 403–408.
- [2] G. B. Djordjević, G. V. Milovanović, *A generalization of class of Panda's polynomials*, VI Conference of Applied Mathematics, Belgrade (1989), 42–47.
- [3] M. Celine Fasenmyer, *Some generalized hypergeometric polynomials*, Bull. Amer. Math. Soc. 53 (1947), 105–108.
- [4] R. Panda, *On new class of polynomials*, Glasgow Math. J. 18 (1977), 105–108.
- [5] E. D. Rainville, *Special functions*, Macmillan Company, New York, 1960.
- [6] Th. M. Rassias, H. M. Srivastava, *Some general families of generating functions for the Laguerre polynomials*, J. Math. Anal. Appl. 174 (1993), 528–538.
- [7] R. C. Singh Chandel, *A note on some generating functions*, Indiana J. Math. 25 (1983), 185–188.
- [8] H. M. Srivastava, *On the product of two Laguerre polynomials*, Bollettino della Unione Matem. Italiano (4) 5 (1972), 1–6.